Heart Weight KMPCLA_HWT_001

Purpose

To evaluate cardiac size using heart weight and body weight.

Experimental Design

- Minimum number of animals : 7M + 7F
- Age at test: Week 60

Equipment

- fine forceps
- surgical scissors
- fine surgical scissor
- kim wipes (tissues) or surgical compress
- laboratory balance
- labelled jar with fixative
- corkplate or wax board
- pins
- jar containing tap water to rinse the tools

Procedure


Methods and procedures used not including center-specific data entry methods.

- 1. Sacrifice the mouse
- 2. Weigh and record total body weight

3. Place mouse on its back and pin the mouse onto board with extended extremities (inner side of hands and foot)

- 4. Wipe or wet the mouse with 70% ethanol to control hair and dander
- 5. Option A (for mice that are undergoing complete necropsy):
 - Proceed with a complete necropsy and tissue collection according to Centre-specific technical SOP including removal of the heart by dissecting the aortic root immediately above the aortic valves and the superior vena cava above the atria
 - Remove adjacent mediastinal fat pads from the excised heart carefully with forceps
 - Empty heart blood by tapping the heart on a kim wipe (absorbent pad) or surgical compress. Repeat until the heart is totally dry

- Weigh the heart, record the weight in the Centre-specific database, and place the heart in fixative
- 6. Option B (for mice that are not undergoing complete necropsy):
 - Open the skin in the ventral midline and in the direction of the extremities and extend cut to hands and feet
 - Open the muscular abdominal wall in the midline and along the lower margin of the rib cage with small forceps
 - Open rib cage by removing the sternum and adjacent ribs
 - Remove the heart by dissecting the aortic root immediately above the aortic valves and the superior vena cava above the atria
 - Remove adjacent mediastinal fat pads from the excised heart carefully with forceps
 - Empty heart blood by tapping the heart on a kim wipe (absorbent pad) or surgical compress. Repeat until the heart is totally dry
 - Weigh the heart and record the weight in the Centre-specific database
 - Discard the heart



Notes

All data are collected at a local workstation in the necropsy room (attached to a digital balance) and uploaded to the Centre-specific pathology data capture system.

Data QC

Mouse weight between 5 grams and 150 grams

IMPC Parameters (+ontology annotations)

Weight (in mg)

Data Analysis, annotation and display (+statistics)

Statistics: ANOVA/Wilcoxon test using normalized heart weights (for body weight) displayed in boxplot

Parameters and Metadata

Equipment manufacturer KMPCLA_HWT_010_001 | v1.0

procedureMetadata

 Req. Analysis: true
 Req. Upload: true
 Is Annotated: false

 Options: Denver Instrument, A & D, Sartorius AG Germany, Mettler Toledo, Ohaus, Radwag,

Tibia length KMPCLA_HWT_002_001 | v1.1

simpleParameter

| Req. Analysis: false | Req. Upload: false | Is Annotated: true |
|----------------------|--------------------|--------------------|
| | | |

Unit Measured: mm

Heart weight normalised against tibia length KMPCLA_HWT_01

3_001 | v1.3 simpleParameter Derivation: div('KMPCLA_HWT_008_001', 'KMPCLA_HWT_002_001')

Equipment ID KMPCLA_HWT_006_001 | v1.0

procedureMetadata

| Req. Analysis: false | Req. Upload: true | Is Annotated: false |
|----------------------|-------------------|---------------------|
| | | |

Heart weight normalised against body weight KMPCLA_HWT_0 12_001 | v1.0

simpleParameter

| Req. Analysis: false | Req. Upload: false | Is Annotated: false |
|----------------------|--------------------|---------------------|
| | | |

Derivation: div('KMPCLA_HWT_008_001', 'KMPCLA_HWT_007_001')

Body weight KMPCLA_HWT_007_001 | v1.3

simpleParameter

Reg. Analysis: false Reg. Upload: true Is Annotated: false

Unit Measured: g

Experimenter ID KMPCLA_HWT_003_001 | v1.0

procedureMetadata

| Req. Analysis: false | Req. Upload: true | Is Annotated: false |
|----------------------|-------------------|---------------------|
| | | |

Date equipment last calibrated KMPCLA_HWT_009_001 | v1.2

procedureMetadata

| Req. Analysis: false | Req. Upload: false | Is Annotated: false |
|----------------------|--------------------|---------------------|
| | | |
| | | |

Heart weight KMPCLA_HWT_008_001 | v1.2

simpleParameter

| Req. Analysis: false | Req. Upload: true | Is Annotated: true |
|----------------------|-------------------|--------------------|
| | | |
| Unit Measured: mg | | |

Equipment model KMPCLA_HWT_011_001 | v1.0

procedureMetadata

Req. Analysis: true Req. Upload: true Is Annotated: false

Options: TE212, MS104S, P-403, Scout Pro SPU123, AB104-S, AV213C, GF-200, Adventurer Pro, TP-114, Adventurer AX223/E, QUINTIX124-1S, PG3001-S, HR-120, AV2101, AV212C, 201-10,

| | | |
|------|------|--|
| | | |
| | | |
| | | |
| | | |
| | | |

Date of sacrifice KMPCLA_HWT_001_001 | v1.0

procedureMetadata

| Req. Analysis: false | Req. Upload: true | Is Annotated: false |
|----------------------|-------------------|---------------------|
| | | |
| | | |

Method of sacrifice KMPCLA_HWT_005_001 | v1.1

procedureMetadata

| Req. Analysis: false | Req. Upload: true | Is Annotated: false |
|----------------------|-------------------|---------------------|
|----------------------|-------------------|---------------------|

Options: Isoflurane overdose, Pentobarb (0.1ml), Ketamine (110mg/kg)/Xylazine (11mg/kg)/ Antisedan (1mg/kg), Cardiac puncture, Ketamine (137mg/kg)/Xylazine (6.6mg/kg), Carbon dioxide, Cervical dislocation, Exsanguination, Ketamine(100mg/kg)/ Xylazine (10mg/kg)/ Antisedan (1mg/kg), Ketamine (100mg/kg)/Xylazine (10mg/kg), None, Anesthetized, Avertin, Ketamine (110mg/kg)/Xylazine (11mg/kg),
